

2172

**MBBS. 2nd Professional Examination-2003.**

Time. 3hours

M.M. 40.

**Forensic Medicine**

Note: All questions are compulsory.

**Part - A**

Q.1. What are the factors, which would help you in estimating the time since death? (10)

Q.2 Differentiate:(05)

- a) Incised from lacerated wound ✓
- b) Suicidal from homicidal cut throat. ✓
- c) Hanging from Strangulation ✓
- d) Dying Declaration from Dying Deposition ✓
- e) Bruise from Abrasion ✓

Q.3 Write short notes on:(05)

- a) Ricochet of bullets ✓
- b) Sudden infant death syndrome
- c) Live Birth
- d) Diatoms
- e) Pugilistic attitude

**Part - B**

Q.4 Describe the signs symptoms and medico-legal aspect of acute alcoholic intoxication.(5) —

Q.5 Describe the duties of Medical Practitioner in case of suspected poisoning.(5) ✓

P.T.O.

2<sup>nd</sup> Prof MBS 2003  
2172 forensic Medicine

Q.6: Each correct answer shall fetch one mark.

R. No. \_\_\_\_\_ Enrollment No. \_\_\_\_\_ Date \_\_\_\_\_

- 1 Bruising of the muscles of neck and larynx with fracture of thyroid cartilage and hyoid bone is seen in death due to  
a) Hanging, b) Throttling, c) Ligature strangulation, d) Traumatic asphyxia.
- 2 The most reliable methods of identification of a living person is  
a) Photography, b) Dactylography, c) Handwriting, d) Anthropometry.
- 3 Greenish color change of putrefication is first seen in the  
a) Right iliac fossa b) Left iliac fossa, c) Back of neck, d) Genital area.
- 4 Blunt trauma will produce incised-like wounds on the  
a) Scalp, b) Palms, c) Neck, d) Lower abdomen.
- 5 Contrecoup injuries are seen in  
a) Liver, b) Heart, c) Brain, d) All of the above.
- 6 On opening the skull the smell can be noted in poisoning by  
a) Phosphorous, b) Arsenic, c) Barbiturate, d) Alcohol.
- 7 Absolute contra indication for gastric lavage is poisoning by  
a) Corrosive poisons, b) Datura Poisoning, c) Phenol, d) HCN
- 8 BAL is useful in the treatment of poisoning due to  
a) Arsenic, b) Endrine, c) Opium, d) Aspirin
- 9 Kleptomania is an irresistible impulse to  
a) Drink alcohol, b) Take a drug of addiction repeatedly, c) Steal articles of little value d) Mutilate animals.
- 10 The drug commonly used for purpose of chronic homicidal poisoning is  
a) Arsenic, b) Barbiturate, c) Aconite, d) Datura.